DIGESTIVE HEALTH CENTER

Medication Information

This applies to patients on anti-platelet, diabetic or iron medications.

Review this medication list for medications you are currently taking.

You may take your usual morning medications (that are not included on the list) on the morning of your procedure. **It is very important to take your chronic pain, high blood pressure, heart and seizure medications that you take daily.**

Medication	***Instructions***
Miralax and other laxatives	Continue taking as you normally do.
Oral Diabetic Medications	Take **half** of your usual dose of diabetic medication in the morning on the day prior to the exam. Do not take your evening dose of diabetic medication on the day prior to the exam. Do not take any diabetic medication on the morning of your exam. If you usually do Accu-Chek readings at home, please do so in the morning prior to coming to your appointment on the day of your exam.
Diabetic Medications (including insulin and insulin pumps)	Contact your doctor that manages your diabetes for further instructions.
Lovenox (enoxaparin)	Do not take your Lovenox injection the morning of your exam.
Coumadin (warfarin) *(Circle One)*	Do Not Hold Hold 3 Days Hold 5 Days
Pradaxa, Xarelto, Eliquis *(Circle One)*	Do Not Hold Hold 1 Day Hold 2 Days Hold 3 Days
Iron (ferrous sulfate)	Stop taking your iron medication **5 days** before your procedure. The day after your procedure, you may resume taking iron supplements.
Aspirin	There is no need to stop taking aspirin before your procedure.
Plavix (clopidogrel), Effient, Brilinta *(Circle One)*	Do Not Hold Hold 5 Days Hold 7 Days
Phentermine Synergy *(Phentermine, Bupropion, and Naltrexone)*	Hold 4 days
Exenatide IR *(Byetta)* Liraglutide *(Victoza/Saxenda)* Semaglutide *(Rybelsus)* Dulaglutide *(Trulicity)* Exenatide ER *(Bydureon)* Semaglutide *(Ozempic/Wegovy)* Tirzapetide *(Mounjaro/Zepbound)*	Clear liquids for 24 hours prior to arrival.
Colonoscopies only: Empagliflozin *(Jardiance)* Dapagliflozin *(Farxiga)* Canagliflozin *(Invokana)* Bexagliflozin *(Brenzavvy)*	Hold 3 days
Colonoscopies only: Ertugliflozin *(Steglatro)*	Hold 4 days

DIGESTIVE HEALTH CENTER
DigestiveHealthCenter.com

1015 S. Hackett Rd., Waterloo, Iowa 50701 **319.234.5990**

Digestive Health Center complies with applicable federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex.
SPANISH : ATENCIÓN: Si habla español, tiene a su disposición servicios gratuitos de asistencia lingüística.
CHINESE : 注意：如果您使用繁體中文, 您可以免費獲得語言援助服務。請致電

Rev. 11-09-26